

Supplementary Table 1. Nutritional status of children aged 0–59 months in 1997

Variable	No. (%)
Weight-for-age (WAZ in 1997)	
Underweight (< -2.0 z-scores)	86 (30.3)
Normal weight (-2.0 to +1.0 z-scores)	187 (65.8)
Risk of overweight (> +1.0 z-scores)	11 (3.9)
Height-for-age (HAZ in 1997)	
Stunted (< -2.0 z-scores)	127 (45.7)
Normal (-2.0 to +3.0 z-scores)	146 (52.5)
Tall (> +3.0 z-scores)	5 (1.8)
Weight-for-height (WHZ in 1997)	
Wasted (< -2.0 z-scores)	30 (10.8)
Normal (-2.0 to +1.0 z-scores)	220 (78.9)
Possible risk of overweight (> +1.0 to +2.0 z-scores)	18 (6.5)
Overweight (> +2.0 to +3.0 z-scores)	4 (1.4)
Obese (> +3.0 z-scores)	7 (2.5)
Body mass index-for-age (BAZ in 1997)	
Wasted (< -2.0 z-scores)	27 (9.6)
Normal (-2.0 to +1.0 z-scores)	212 (75.2)
Possible risk of overweight (> +1.0 to +2.0 z-scores)	27 (9.6)
Overweight (> +2.0 to +3.0 z-scores)	9 (3.2)
Obese (> +3.0 z-scores)	7 (2.5)

WAZ, weight-for-age z-score; HAZ, height-for-age z-score; WHZ, weight-for-height z-score.